CUMULATIVE AUTHOR INDEX
Environmental Entomology
Volume 28, 1999

A

Ahern, R. G., 681
Akbulut, S., 407, 1014
Alborn, D. H., 973
Allen, W. W., 594
Allison, D., 61
Alyokhin, A. V., 425
Ampong-Nyarko, K., 1154
Andow, D. A., 1092
Appel, A. G., 935, 979
Arbogast, R. T., 557
Armer, C. A., 1036
Arn, H., 947
Arthur, F. H., 529
Ashikaga, T., 1085

B

Baker, J. E., 529
Balciunas, J. K., 290
Barnard, D. R., 545
Barrows, E. M., 795
Baxendale, F. P., 123
Beanland, L., 1101
Beckage, N., 536
Beggs, J. R., 9
Beisner, B. E., 1107
Bell, K. L., 163
Bellows, T. S., 282, 300
Bengtsson, M., 775
Bentz, B. J., 577, 924
Bentz, J.-A., 456
Berisford, C. W., 638, 649
Biddick, L. L., 954
Binns, M., 398
Bitzer, R. J., 608
Blatt, S. E., 246
Bledsoe, L. W., 188
Bloem, K. A., 669
Bloem, S., 669
Booth, G. M., 812
Borden, J. H., 246
Bossart, J. L., 628
Bostanian, N. J., 398
Boydston, L., 61
Brewer, G. J., 1172
Briano, J. A., 88
Briere, J.-F., 22
Brinkman, M. A., 863
Brodeur, J., 622, 735
Brown, L. G., 895
Burkness, E. C., 851
Burrows, D. W., 290
Buschini, M. L. T., 618
Buss, L. J., 485
Butler, L., 795

C

Calcaterra, L. A., 88
Calkins, C. O., 669
Carlson, D. A., 545
Carmona, D. M., 1145
Carpenter, J. E., 551, 669
Carr, T. G., 461
Cartagena, M. C., 420
Casey, C. A., 998
Chihrane, J., 116
Chilcutt, C. F., 505
Cloutier, C., 735
Consolie, N., 890
Cook, C. A., 173
Cooke, B. J., 711
Copren, K. A., 1077
Cornelius, M. L., 1140
Crist, T. O., 659, 681
Croft, B. A., 108, 322, 523, 915
Cronin, J. T., 496
Croughan, T. P., 1130
Czesak, M. E., 217

D

Dadour, I. R., 675
Dahlsten, D. L., 845
Dalthorp, D., 255
Dalusky, M. J., 638
Darnell, S. J., 266
Debouzie, D., 53
De Farias, A. M. I., 858
Degooyer, T. A., 703
de Mackiewicz, D., 470
Di Fonzo, C. D., 188
Doğan, E. B., 812
Doostdar, H., 353
Downie, D. A., 1004
Dreistadt, S. H., 845
Duan, J. J., 1140

E

Eckel, R. V. W., 566
Edelson, J. V., 572
Edwards, C. R., 188
Ekesi, S., 1154
Elkinton, J. S., 240
Ellington, J., 722
Elliott, N. C., 597
El-Sayed, A., 775, 947
English-Loeb, G., 890
Evans, T. A., 823

F

Facundo, H. T., 14
Ferro, D. N., 425
Fisher, J. R., 347
Flanders, K. L., 128
Flechtmann, C. A. H., 638, 649
Fleischer, S. J., 470
Flinn, P. W., 529
Forsline, A., 890
Fox, C. W., 217
Freitas, S., 178
French, B. W., 597
Fuller, B. W., 863
Fuxa, J. R., 330

G

Galante, E., 420
Garrido, A., 339
Geden, C. J., 755
Getty, G. M., 1077
Gildow, F. E., 470
Gillespie, R., 61
Gleeson, P. V., 823
Gödde, J., 947
Gómez-Guillamón, M. L., 831

Gotway, C. A., 266
Gouge, D. H., 876
Gouli, S., 1085
Graf, G., 61
Granett, J., 1004
Greco, A. M., 1044
Gruenhagen, N. M., 884

H
Haile, F. J., 787, 1136
Hajek, A. E., 845
Hamilton, R. M., 812
Hammack, L., 961
Harmsen, R., 72
Harris, A., 275
Harris, M. K., 572
Haverty, M. I., 1077
Hawthorne, D. J., 696
Hayes, J. L., 496
Headrick, D. H., 282, 300, 537
Heagle, A. S., 566
Heng-Moss, T. M., 123
Henneberry, T. J., 876
Herbers, J. M., 538
Hibbard, B. E., 961
Higley, L. G., 787, 1136
Hoback, W. W., 1136
Hofstetter, R. W., 954
Holst, N., 836, 1060
Holyoke, C. W., 961
Hopper, K. R., 858
Horgan, F. G., 96
Hoy, C. W., 1101
Hunt, T. E., 1136
Hunter, W. B., 895
Hutchison, W. D., 445, 851

I
Inbar, M., 353
Irwin, M. E., 1036
Isard, S. A., 188
Ishihara, M., 195
Ishikawa, Y., 690

J
Jacas, J.-A., 339
Jackson, D. M., 566
James, R. R., 108
Jensen, T., 545
Jewett, D. K., 551
Johnson, S. J., 330
Joselyn, M. G., 188
Joyce, A. L., 282
Jyoti, J. L., 1172

K
Katagiri, C., 968
Keever, G. J., 935
Kemp, W. P., 347
Kendra, P. E., 557
Kimura, M. T., 968
King, E. G., 137
Kline, M., 961
Knight, A. L., 81
Knighten, K., 275
Kondo, V., 795
Kopanic, R. J., Jr., 431
Kovach, J., 398
Kukan, B., 44

L
Lacey, L. A., 913
Lacourse, C. D., 1044
Landis, D. A., 1145
Landolt, P. J., 954, 973
Landry, J. B., 1044
Lapointe, S. L., 895
Laugé, G., 116
Lee, J.-H., 1177
Lee, L. L., 876
Lenz, M., 823
Leonard, D. E., 240
Le Roux, A.-Y., 22
Lester, P. J., 72
Leva, D. M., 961
Levine, E., 188
Lewis, C. N., 986
Lewis, E. E., 907
Lewis, V. R., 1077
Lill, J. T., 1114
Lim, U. T., 1177
Linit, M. J., 407, 1014
Linn, C. E., Jr., 14
Liu, T.-X., 901
Llácer, E., 339
Löfqvist, J., 775
Logan, J. A., 143, 924
López-Arroyo, J. I., 1183
López-Sesé, A. I., 831
Lucas, E., 622
Luh, H. K., 915
Lukezic, F. L., 470
Luttrell, R. G., 275
Lymbery, A. J., 675
Lysyk, T. J., 387

M
Mack, T. P., 935
Mailloux, G., 398
Maisonhaute, C., 116
Malham, J. P., 9
Maniania, N. K., 1154
Mankin, R. W., 557
Markham, R. H., 836
Marshall, S. D., 1052
Martinson, T., 890
Mastro, V. C., 240
Mathews, P. L., 729
Matter, S. F., 1044
Mayer, R. T., 353
Mazih, A., 53
McCullough, D. G., 485
McPherson, R. M., 768
Meagher, R. L., Jr., 367
Meikle, W. G., 836
Meinke, L. J., 266
Melo, R., 513
Merritt, R. W., 233
Messing, R. H., 1140
Michaud, J. P., 205
Miller, J. C., 518
Miller, J. M., 128
Miller, R., 61
Miller, S. A., 1101
Miller, T., 61
Mills, N., 748
Mills, N. J., 868
Mitchell, E. R., 367
Moar, W. J., 979
Morales-Ramos, J. A., 137
Muegge, M. A., 1130
Mulla, M. S., 761
Mullins, D. E., 577
Myers, J. H., 44, 96, 1107

N
Nault, L. R., 1101
Neal, J. J., 173
Ni, X., 787
Nielsen, P. S., 1060

Nyrop, J., 255
Nyrop, J. P., 1021

O

Ohtsu, T., 968
Oliveira, L., 513
Onstad, D. W., 188
Onu, I., 1154
O'Rourke, P. K., 851
Osborne, K. H., 594
Ottati, A. L. T., 649

P

Parker, B. L., 1085
Parker, C. D., Jr., 275
Parker, P. E., 315
Päts, P., 1124
Pearson, G. A., 943
Peck, D. C., 372
Pedigo, L. P., 703
Peng, F., 330
Perez-Mendoza, J., 529
Perring, T. M., 300, 884
Pickett, J. A., 983
Pierre, J.-S., 22
Pierson, F. B., 347
Pike, K. S., 61
Pleasants, J. M., 608
Poprawski, T. J., 315
Portillo-Aguilar, C., 1021
Powell, J. S., 148
Pracros, P., 22
Pree, D. J., 72
Purcell, A. H., 868

Q

Quisenberry, S. S., 477, 787

R

Rabbinge, R., 307
Racette, G., 398
Radford, B. J., 163
Raffa, K. F., 148
Ramaswamy, S. B., 275
Ramm, C. W., 485
Raupp, M. J., 998
Rauscher, S., 775
Reed, J., 275
Régnière, J., 711
Rice, M. E., 703
Rice, W. C., 1130
Richter, A. R., 330
Ring, D. R., 1130
Riordan, T. P., 123
Robbins, J., 275
Robertson, L. N., 163
Rodriguez-Saona, C., 518
Roelofs, W. L., 14
Roininen, H., 461
Rojas, J. C., 155, 588
Rojas, M. G., 137
Rothwell, N. L., 358
Rowney, D. L., 845
Roy, M., 735
Royer, T. A., 572
Rypstra, A. L., 1052

S

Salvo, A., 539
Sanchez-Martinez, G., 38
Sárospataki, M., 1
Savalli, U. M., 217
Schaalje, G. B., 812
Schal, C., 431, 943
Schausberger, P., 322, 523
Schellhorn, N. A., 1092
Schmaedick, M. A., 439
Schotzko, D. J., 477
Scriber, J. M., 628
Shah, P. A., 983
Shapiro, D. I., 907
Shelly, T. E., 780
Shelton, A. M., 439
Shields, E. J., 128
Shimada, M., 195
Shintani, Y., 690
Simmons, A. M., 212
Simpson, G. B., 163
Skinner, M., 1085
Skovgård, H., 1060
Smith, L. M., II, 935
Smitley, D. R., 358
Soria, C., 831
Southward, M., 722
Spence, J. R., 224
Spencer, J. L., 188
Stansly, P. A., 901
Stark, D. M., 868
Starý, P., 61
Steiner, C. A., 496
Stephen, F. M., 729
Stewart, S., 275
Stout, M. J., 1130
Strauss, S. H., 108
Strickler, K., 178, 1067
Su, T., 761
Sutula, C., 275

T

Tabashnik, B. E., 505
Tang, Y. Q., 895
Tanley, M. J., 979
Tauber, C. A., 1021, 1183
Tauber, M. J., 1021, 1183
Tavares, J., 513
Teillon, H. B., 1085
Testa, A., 128
Thaler, J. S., 30
Thistlewood, H. M. A., 72
Tikkanen, O.-P., 461
Toft, R. J., 9
Tomás, O., 339
Townsend, A. M., 456
Townsend, E. C., 795
Trevisan, L. M., 72
Tsai, J. H., 315, 412
Tumlinson, J. H., 973
Turchin, P., 496
Turner, J. E., 81

U

Urbaneja, A., 339

V

Valladares, G. R., 539
Vandenberg, J. D., 983
Van Der Werf, W., 307
Van Meel, R., 96
Vega, F. E., 535, 913
Venette, R. C., 445, 851
Vernon, R. S., 1124
Villani, M., 255, 890
Villani, M. G., 14, 1021
Volney, W. J. A., 224

W

Wagner, M. R., 38
Wallace, J. R., 233
Wang, K., 412
Weaver, D. K., 557
Weber, J. D., 224

Wells, M. L., 768
Weseloh, R. M., 1162
Whitbey, R. M., 201
Whitfield, J. B., 912, 986
Wiedenmann, R. N., 1036
Williams, D. F., 88
Williams, J. A., 659
Willson, H., 188
Wilson-Rummenie, A. C., 163
Witzgall, P., 775

X

Xia, J., 275
Xia, J. Y., 307

Y

Young, L. J., 123, 266

Z

Zack, R. S., 536
Zaviezo, T., 748
Zeng, F., 275
Zlotina, M. A., 240
Zuparko, R. L., 845

CUMULATIVE SUBJECT INDEX
Environmental Entomology
Volume 28, 1999

A

above-below-ground growth,
compensatory growth, 38
Neodiprion gillettei, 38
Pinus ponderosa, 38
sawfly defoliation, 38
abundance,
Macrolepidoptera, 795
sampling, 795
species richness, 795
traps, 795
weather, 795
Acalymma vittatum,
bacterial wilt, 470
cucurbits, 470
epidemiology, 470
Erwinia tracheiphila, 470
acclimatization,
heat shock, 116
heat shock protein persistence, 116
thermotolerance, 116
Trichogramma brassicae, 116
Adelges tsugae,
insect coldhardiness, 1085
Tsuga canadensis, 1085
Tsuga caroliniana, 1085
adult abundance,
Anthonomus signatus, 398
management, 398
predictive model, 398
Ageneotettix deorum,
grasshoppers, 347
hatch prediction, 347
models, 347
postdiapause development, 347
rangeland, 347
aggregation,
peas, 477
Sitona lineatus, 477
spatiotemporal analysis, 477
tillage, 477
alarm pheromone,
aphid, 983
Diuraphis noxia, 983
Myzus persicae, 983
Paecilomyces fumosoroseus, 983
semiochemicals, 983
alfalfa,
foraging behavior, 1067
Medicago sativa, 1067
Megachile rotundata, 1067
pollination ecology, 1067
seed production, 1067
alfalfa snout beetle,
entomopathogenic nematode, 128
Heterorhabditis bacteriophora, 128
Amblyseius fallacis,
biological control, 72
encapsulated cypermethrin, 72
Panonychus ulmi, 72
Anthonomus grandis,
Callosobruchus maculatus, 137
Catolaccus grandis, 137
factitious host, 137
host acceptance, 137
Anthonomus signatus,
adult abundance, 398
management, 398
predictive model, 398
antibiosis,
chlorophyll fluorescence, 787
photosynthesis, 787
Russian wheat aphid, 787
tolerance, 787
wheat, 787
Aphelinus asychis,
Aphidius matricariae, 858
biological control, 858
Diuraphis noxia, 858
host acceptance, 858
host defense, 858
aphid,
alarm pheromone, 983
Diuraphis noxia, 983
Myzus persicae, 983
Paecilomyces fumosoroseus, 983
semiochemicals, 983
aphid parasitoid,
Diaeretiella rapae, 61
habitats, 61
host range, 61
host switching, 61

Aphidius matricariae,
Aphelinus asychis, 858
biological control, 858
Diuraphis noxia, 858
host acceptance, 858
host defense, 858
Apis,
Bombus, 367
Cerceris, 367
Larra, 367
nontarget insects, 367
Apis mellifera,
Bacillus thuringiensis, 1172
Cochylis hospes, 1172
Helianthus annuus, 1172
apple,
leafrollers, 81
mating disruption, 81
Pandemis, 81
sex pheromones, 81
argan fruit,
Ceratitis capitata, 53
mortality factors, 53
number of generations, 53
oviposition, 53
population dynamics, 53
arrestant,
attractant, 954
Cydia pomonella, 954
deterrent, 954
repellent, 954
arthropod,
developmental rate, 22
insect, 22
model, 22
temperature, 22
Asclepias,
florivory, 1044
flowers, 1044
herbivory, 1044
Tetraopes, 1044
aster yellows phytoplasma,
gender, 1101
lettuce, 1101
Macrosteles quadrilineatus, 1101
transmission, 1101
vector, 1101
Ataenius spretulus,
Bacillus popilliae, 358
Lolium perenne, 358
mowing height, 358
predators, 358
turfgrass, 358
attractant,
arrestant, 954
Cydia pomonella, 954
deterrent, 954
repellent, 954
attractants,
Diabrotica barberi, 961
Diabrotica virgifera virgifera, 961
maize silk volatiles, 961
repellents, 961
semiochemicals, 961
attraction,
Colorado potato beetle, 973
host-finding, 973
Leptinotarsa decemlineata, 973
volicitin, 973
Aulacidea pilosellae,
biological control, 1
Hieracium pilosella, 1
phytophagous insect fauna, 1
Tephritis ruralis, 1
azalea,
azalea lace bug, 998
development time, 998
mesophyll-feeder, 998
nitrogen fertilizer, 998
survivorship, 998
azalea lace bug,
azalea, 998
development time, 998
mesophyll-feeder, 998
nitrogen fertilizer, 998
survivorship, 998

B

Bacillus popilliae,
Ataenius spretulus, 358
Lolium perenne, 358
mowing height, 358
predators, 358
turfgrass, 358
Bacillus thuringiensis, 108, 1172
Apis mellifera, 1172
Chrysomela scripta, 108
Cochylis hospes, 1172
cottonwood, 108
genetic engineering, 108
Helianthus annuus, 1172
insect resistance, 108
poplars, 108

bacterial wilt,
Acalymma vittatum, 470
cucurbits, 470
epidemiology, 470
Erwinia tracheiphila, 470
Bactrocera dorsalis,
olfactory stimuli, 1140
pest sampling, 1140
protein, 1140
visual stimuli, 1140
baculovirus,
enzyme-linked immunosorbent assay, 868
epizootiology, 868
grape, 868
microbial control, 868
vineyard, 868
bait,
Blattella germanica, 431
coprophagy, 431
horizontal toxicant transfer, 431
bark beetles,
Dendroctonus frontalis, 496
dispersal, 496
mark-recapture experiment, 496
pest management strategies, 496
spatial dynamics, 496
Beauveria bassiana,
natural enemies, 863
rangeland grasshoppers, 863
behavior,
Coenosia tigrina, 1124
Delia radicum, 1124
exclusion fences, 1124
insect pest management, 1124
physical control, 1124
Bemisia argentifolii,
induced response, 353
interspecific competition, 353
omnivory, 353
Trichoplusia ni, 353
Bemisia tabaci,
life history parameters, 831
melon, 831
resistance, 831
whitefly, 831
binomial sampling,
bootstrap simulation, 851
carrot, 851
Macrosteles quadrilineatus, 851
resampling software, 851
biodiversity,
Braconidae, 986
complementarity, 986
forest management, 986
species richness, 986
biological control, 1, 72, 88, 96, 529, 858, 890, 901, 915, 1145, 1154, 1177, 1183
Amblyseius fallacis, 72
Aphelinus asychis, 858
Aphidius matricariae, 858
Aulacidea pilosellae, 1
Ceraeochrysa spp., 1183
Coccinellid, 901
colony composition, 88
conservation, 1145
cowpea, 1154
Cyzenis albicans, 96
Daktulosphaira vitifolia, 890
development, 1183
diatomaceous earth, 529
discriminant analysis, 915
Diuraphis noxia, 858
encapsulated cypermethrin, 72
entomopathogenic nematodes, 890
enzyme-linked immunosorbent assay, 1177
feeding behavior, 901
fire ants, 88
generalist-specialist predators, 915
genetic algorithm, 915
ground beetles, 1145
habitat management, 1145
Hieracium pilosella, 1
host acceptance, 858
host defense, 858
inert dust, 529
lady beetle, 901
mass rearing, 1183
Megalurothrips sjostedti, 1154
Metarhizium anisopliae, 1154
Nilaparvata lugens, 1177
Operophtera brumata, 96
Panonychus ulmi, 72
parasitic ants, 88
parasitoid, 529
phylloxera, 890
phytophagous insect fauna, 1
pitfall traps, 1145
predation, 1177
predators, 901

refuge habitats, 1145
reproduction, 1183
simultaneous introduction, 96
soil moisture, 890
Solenopsis daguerrei, 88
Solenopsis richteri, 88
spider, 1177
spider mites, 915
stored products, 529
synthetic insecticide nontarget organisms, 1154
Tephritis ruralis, 1
Tetranychidae, 915
thrips, 1154
Vitis, 890
whitefly, 901
winter moth, 96
biological invasions,
climate matching software, 445
cotton, 445
establishment, 445
Pectinophora gossypiella, 445
risk assessment, 445
Blattella germanica,
bait, 431
coprophagy, 431
horizontal toxicant transfer, 431
body water content,
Callosobruchus maculatus, 979
cuticular permeability, 979
desiccants, 979
insecticidal dusts, 979
water loss, 979
Bombus,
Apis, 367
Cerceris, 367
Larra, 367
nontarget insects, 367
bootstrap simulation,
binomial sampling, 851
carrot, 851
Macrosteles quadrilineatus, 851
resampling software, 851
Braconidae,
biodiversity, 986
complementarity, 986
forest management, 986
species richness, 986
brown citrus aphid,
citrus tristeza, 895
life table, 895
population growth, 895
vector, 895
virus, 895
buffalograss,
mealybug, 123
parasitoid, 123
Rhopus nigroclavatus, 123
Tridiscus sporoboli, 123
Bursaphelenchus xylophilus, 407, 1014
fertility table, 407
flight, 1014
flight mill, 1014
intrinsic rate of increase, 407
Monochamus carolinensis, 407
pine wilt disease, 1014
pinewood nematode, 1014

C

cadaver,
Heterorhabditis bacteriophora, 907
infectivity, 907
Steinernema carpocapsae, 907
Callosobruchus maculatus, 137, 979
Anthonomus grandis, 137
body water content, 979
Catolaccus grandis, 137
cuticular permeability, 979
desiccants, 979
factitious host, 137
host acceptance, 137
insecticidal dusts, 979
water loss, 979
cannibalism, 523, 823, 1092
coccinellids, 1092
Coptotermes acinaciformis, 823
dispersal, 1092
interspecific intraguild predation, 1092
interspecific predation, 523
mark–recapture, 823
mound-building, 823
Nile Blue A, 823
predaceous mites, 523
predation preference, 523
prey scarcity, 1092
canopy distribution,
Eucallipterus tiliae, 845
Trioxys curvicaudus, 845
carrot,
binomial sampling, 851
bootstrap simulation, 851
Macrosteles quadrilineatus, 851

resampling software, 851
caterpillars,
experimental epizootiology, 1107
Malacosoma californicum pluviale, 1107
nuclearpolyhedrovirus, 1107
population cycles, 1107
virus transmission, 1107
Catolaccus grandis,
Anthonomus grandis, 137
Callosobruchus maculatus, 137
factitious host, 137
host acceptance, 137
Ceraeochrysa spp.,
biological control, 1183
development, 1183
mass rearing, 1183
reproduction, 1183
Ceratitis capitata, 53, 780
argan fruit, 53
mating behavior, 780
mortality factors, 53
number of generations, 53
oviposition, 53
population dynamics, 53
trimedlure, 780
Cerceris,
Apis, 367
Bombus, 367
Larra, 367
nontarget insects, 367
chlorophyll fluorescence,
antibiosis, 787
photosynthesis, 787
Russian wheat aphid, 787
tolerance, 787
wheat, 787
Chrysomela scripta,
Bacillus thuringiensis, 108
cottonwood, 108
genetic engineering, 108
insect resistance, 108
poplars, 108
citrus tristeza,
brown citrus aphid, 895
life table, 895
population growth, 895
vector, 895
virus, 895
climate change,
direct temperature control, 924
mountain pine beetle, 924
phenology, 924
seasonality, 924
temperature-dependent development, 924
climate matching software,
biological invasions, 445
cotton, 445
establishment, 445
Pectinophora gossypiella, 445
risk assessment, 445
Coccinellid,
biological control, 901
feeding behavior, 901
lady beetle, 901
predators, 901
whitefly, 901
coccinellids,
cannibalism, 1092
dispersal, 1092
interspecific intraguild predation, 1092
prey scarcity, 1092
Cochylis hospes,
Apis mellifera, 1172
Bacillus thuringiensis, 1172
Helianthus annuus, 1172
Coenosia tigrina,
behavior, 1124
Delia radicum, 1124
exclusion fences, 1124
insect pest management, 1124
physical control, 1124
cold heat tolerance,
Drosophila, 968
energy storage, 968
phospholipids, 968
triacylglycerols, 968
colony composition,
biological control, 88
fire ants, 88
parasitic ants, 88
Solenopsis daguerrei, 88
Solenopsis richteri, 88
Colorado potato beetle, 425, 973
attraction, 973
flight, 425
host-finding, 973
Leptinotarsa decemlineata, 973
mating, 425
resistance management, 425

transgenic plants, 425
volicitin, 973
community structure,
density dependence, 1114
Diadegma psilocorsis, 1114
parasitoids, 1114
Psilocorsis quercicella, 1114
Quercus alba, 1114
compensatory growth,
above-below-ground growth, 38
Neodiprion gillettei, 38
Pinus ponderosa, 38
sawfly defoliation, 38
complementarity,
biodiversity, 986
Braconidae, 986
forest management, 986
species richness, 986
computer simulation,
integrated pest management, 505
population models, 505
conservation,
biological control, 1145
ground beetles, 1145
habitat management, 1145
pitfall traps, 1145
refuge habitats, 1145
conservation biological control,
facultative phytophagy, 1036
flower bug, 1036
insect–plant interactions, 1036
Orius insidiosus, 1036
coprophagy,
bait, 431
Blattella germanica, 431
horizontal toxicant transfer, 431
Coptotermes acinaciformis,
cannibalism, 823
mark–recapture, 823
mound-building, 823
Nile Blue A, 823
cotton,
biological invasions, 445
climate matching software, 445
establishment, 445
Pectinophora gossypiella, 445
risk assessment, 445
cottonwood,
Bacillus thuringiensis, 108
Chrysomela scripta, 108
genetic engineering, 108
insect resistance, 108
poplars, 108
cowpea,
biological control, 1154
Megalurothrips sjostedti, 1154
Metarhizium anisopliae, 1154
synthetic insecticide nontarget organisms, 1154
thrips, 1154
cucurbits,
Acalymma vittatum, 470
bacterial wilt, 470
epidemiology, 470
Erwinia tracheiphila, 470
cuticular permeability,
body water content, 979
Callosobruchus maculatus, 979
desiccants, 979
insecticidal dusts, 979
water loss, 979
Cydia pomonella, 775, 954
(E, E)-8,10-dodecadienol, 775
arrestant, 954
attractant, 954
deterrent, 954
Lepidoptera, 775
repellent, 954
sex pheromone, 775
Tortricidae, 775
wind tunnel, 775
Cyzenis albicans,
biological control, 96
Operophtera brumata, 96
simultaneous introduction, 96
winter moth, 96

D

Daktolosphaira vitifoliae,
distribution, 1004
extinction-colonization, 1004
grape phylloxera, 1004
population size, 1004
Vitis, 1004
Daktulosphaira vitifolia,
biological control, 890
entomopathogenic nematodes, 890
phylloxera, 890
soil moisture, 890
Vitis, 890
defoliation,
economic injury level, 1136
Glycine max, 1136

herbivory, 1136
insect injury, 1136
leaf area index, 1136
degree-days,
developmental threshold, 518
Hippodamia convergens, 518
mortality, 518
weights, 518
dehydration,
mortality, 935
Periplaneta americana, 935
Periplaneta fuliginosa, 935
saturation deficit, 935
water relations, 935
delayed flood,
herbicide resistant germplasm, 1130
Lissorhoptrus oryzophilus, 1130
rice plant, 1130
rice water weevil, 1130
Delia radicum,
behavior, 1124
Coenosia tigrina, 1124
exclusion fences, 1124
insect pest management, 1124
physical control, 1124
Dendroctonus frontalis,
bark beetles, 496
dispersal, 496
mark-recapture experiment, 496
pest management strategies, 496
spatial dynamics, 496
density dependence,
community structure, 1114
Diadegma psilocorsis, 1114
parasitoids, 1114
Psilocorsis quercicella, 1114
Quercus alba, 1114
desiccants,
body water content, 979
Callosobruchus maculatus, 979
cuticular permeability, 979
insecticidal dusts, 979
water loss, 979
deterrent,
arrestant, 954
attractant, 954
Cydia pomonella, 954
repellent, 954
development, 513, 1183
biological control, 1183
Ceraeochrysa spp., 1183
Glyptapanteles militaris, 513
host age, 513
mass rearing, 1183
parasitoid, 513
Pseudaletia unipuncta, 513
reproduction, 1183
development time,
azalea, 998
azalea lace bug, 998
mesophyll-feeder, 998
nitrogen fertilizer, 998
survivorship, 998
developmental rate, 22, 412
arthropod, 22
insect, 22
life table, 412
model, 22
temperature, 22
Toxoptera citricida, 412
developmental threshold,
degree-days, 518
Hippodamia convergens, 518
mortality, 518
weights, 518
Diabrotica barberi,
attractants, 961
Diabrotica virgifera virgifera, 961
maize silk volatiles, 961
repellents, 961
semiochemicals, 961
Diabrotica virgifera virgifera,
attractants, 961
Diabrotica barberi, 961
maize silk volatiles, 961
repellents, 961
semiochemicals, 961
Diadegma psilocorsis,
community structure, 1114
density dependence, 1114
parasitoids, 1114
Psilocorsis quercicella, 1114
Quercus alba, 1114
Diaeretiella rapae,
aphid parasitoid, 61
habitats, 61
host range, 61
host switching, 61
diapause,
Haematobia irritans, 387
simulation model, 387
spring emergence, 387

temperature, 387
diatomaceous earth,
biological control, 529
inert dust, 529
parasitoid, 529
stored products, 529
direct temperature control,
climate change, 924
mountain pine beetle, 924
phenology, 924
seasonality, 924
temperature-dependent development, 924
discriminant analysis,
biological control, 915
generalist-specialist predators, 915
genetic algorithm, 915
spider mites, 915
Tetranychidae, 915
dispersal, 372, 496, 1092
bark beetles, 496
cannibalism, 1092
coccinellids, 1092
Dendroctonus frontalis, 496
interspecific intraguild predation, 1092
mark-recapture experiment, 496
pest management strategies, 496
population dynamics, 372
prey scarcity, 1092
Prosapia nr *bicincta*, 372
Prosapia simulans, 372
spatial dynamics, 496
distribution,
Daktolosphaira vitifoliae, 1004
extinction-colonization, 1004
grape phylloxera, 1004
population size, 1004
Vitis, 1004
Diuraphis noxia, 858, 983
alarm pheromone, 983
Aphelinus asychis, 858
aphid, 983
Aphidius matricariae, 858
biological control, 858
host acceptance, 858
host defense, 858
Myzus persicae, 983
Paecilomyces fumosoroseus, 983
semiochemicals, 983
Drosophila,
cold heat tolerance, 968
energy storage, 968
phospholipids, 968
triacylglycerols, 968
dung beetle communities,
large herbivorous dung, 420
Mediterranean ecosystem, 420
rabbit dung, 420
Scarabaeoidea, 420
trophic resources, 420

E

(E, E)-8,10-dodecadienol,
Cydia pomonella, 775
Lepidoptera, 775
sex pheromone, 775
Tortricidae, 775
wind tunnel, 775
economic injury level,
defoliation, 1136
Glycine max, 1136
herbivory, 1136
insect injury, 1136
leaf area index, 1136
emergence patterns,
mortality, 9
parasitoid, 9
Shecophaga, 9
Vespula, 9
yellowjackets, 9
encapsulated cypermethrin,
Amblyseius fallacis, 72
biological control, 72
Panonychus ulmi, 72
energy storage,
cold heat tolerance, 968
Drosophila, 968
phospholipids, 968
triacylglycerols, 968
entomopathogenic nematode,
alfalfa snout beetle, 128
Heterorhabditis bacteriophora, 128
entomopathogenic nematodes, 876, 890
biological control, 890
Daktulosphaira vitifolia, 890
Heterorhabditis, 876
infectivity, 876
lepidopteran cotton pests, 876
phylloxera, 890
soil moisture, 890
Steinernema, 876

temperature, 876
Vitis, 890
Entomophaga maimaiga, 485, 1162
100-tree plots, 485
fixed-radius plots, 485
infection model, 1162
infection reservoir, 1162
Lymantria dispar, 485, 1162
nuclear polyhedrosis virus, 485
rainfall, 1162
resting spores, 1162
timed walks, 485
enzyme-linked immunosorbent assay, 868, 1177
baculovirus, 868
biological control, 1177
epizootiology, 868
grape, 868
microbial control, 868
Nilaparvata lugens, 1177
predation, 1177
spider, 1177
vineyard, 868
Ephestia kuehniella,
flour mills, 1060
integrated control, 1060
population dynamics, 1060
simulation model, 1060
temperature, 1060
epidemiology,
Acalymma vittatum, 470
bacterial wilt, 470
cucurbits, 470
Erwinia tracheiphila, 470
epizootiology,
baculovirus, 868
enzyme-linked immunosorbent assay, 868
grape, 868
microbial control, 868
vineyard, 868
Erwinia tracheiphila,
Acalymma vittatum, 470
bacterial wilt, 470
cucurbits, 470
epidemiology, 470
establishment,
biological invasions, 445
climate matching software, 445
cotton, 445
Pectinophora gossypiella, 445
risk assessment, 445
Eucallipterus tiliae,
canopy distribution, 845
Trioxys curvicaudus, 845
exclusion fences,
behavior, 1124
Coenosia tigrina, 1124
Delia radicum, 1124
insect pest management, 1124
physical control, 1124
Exomala orientalis,
golf course, 14
temporal spatial distribution, 14
experimental epizootiology,
caterpillars, 1107
Malacosoma californicum pluviale, 1107
nuclearpolyhedrovirus, 1107
population cycles, 1107
virus transmission, 1107
extinction-colonization,
Daktolosphaira vitifoliae, 1004
distribution, 1004
grape phylloxera, 1004
population size, 1004
Vitis, 1004

F

factitious host,
Anthonomus grandis, 137
Callosobruchus maculatus, 137
Catolaccus grandis, 137
host acceptance, 137
facultative phytophagy,
conservation biological control, 1036
flower bug, 1036
insect–plant interactions, 1036
Orius insidiosus, 1036
feeding behavior, 901, 1077
biological control, 901
Coccinellid, 901
foraging behavior, 1077
lady beetle, 901
predators, 901
subterranean termites, 1077
whitefly, 901
wood consumption, 1077
feeding injury,
Freeman maple, 456
nymphal survivorship, 456
oviposition, 456

potato leafhopper, 456
red maple, 456
female behavior,
pheromone, 943
pheromone autodetection, 943
Vitacea polistiformis, 943
fertility table,
Bursaphelenchus xylophilus, 407
intrinsic rate of increase, 407
Monochamus carolinensis, 407
fire ants,
biological control, 88
colony composition, 88
parasitic ants, 88
Solenopsis daguerrei, 88
Solenopsis richteri, 88
fixed-radius plots,
100-tree plots, 485
Entomophaga maimaiga, 485
Lymantria dispar, 485
nuclear polyhedrosis virus, 485
timed walks, 485
flight, 425, 1014
Bursaphelenchus xylophilus, 1014
Colorado potato beetle, 425
flight mill, 1014
mating, 425
pine wilt disease, 1014
pinewood nematode, 1014
resistance management, 425
transgenic plants, 425
flight mill,
Bursaphelenchus xylophilus, 1014
flight, 1014
pine wilt disease, 1014
pinewood nematode, 1014
florivory,
Asclepias, 1044
flowers, 1044
herbivory, 1044
Tetraopes, 1044
flour mills,
Ephestia kuehniella, 1060
integrated control, 1060
population dynamics, 1060
simulation model, 1060
temperature, 1060
flower bug,
conservation biological control, 1036
facultative phytophagy, 1036
insect–plant interactions, 1036
Orius insidiosus, 1036
flowers,
Asclepias, 1044
florivory, 1044
herbivory, 1044
Tetraopes, 1044
foraging behavior, 1067, 1077
alfalfa, 1067
feeding behavior, 1077
Medicago sativa, 1067
Megachile rotundata, 1067
pollination ecology, 1067
seed production, 1067
subterranean termites, 1077
wood consumption, 1077
forest Lepidoptera,
Malacosoma californicum pluviale, 44
nuclear polyhedrosis virus, 44
population cycles, 44
population dynamics, 44
tent caterpillars, 44
forest management,
biodiversity, 986
Braconidae, 986
complementarity, 986
species richness, 986
Freeman maple,
feeding injury, 456
nymphal survivorship, 456
oviposition, 456
potato leafhopper, 456
red maple, 456

G

gender,
aster yellows phytoplasma, 1101
lettuce, 1101
Macrosteles quadrilineatus, 1101
transmission, 1101
vector, 1101
generalist-specialist predators,
biological control, 915
discriminant analysis, 915
genetic algorithm, 915
spider mites, 915
Tetranychidae, 915
genetic algorithm,
biological control, 915
discriminant analysis, 915
generalist-specialist predators, 915

spider mites, 915
Tetranychidae, 915
genetic engineering,
Bacillus thuringiensis, 108
Chrysomela scripta, 108
cottonwood, 108
insect resistance, 108
poplars, 108
Glycine max,
defoliation, 1136
economic injury level, 1136
herbivory, 1136
insect injury, 1136
leaf area index, 1136
Glyptapanteles militaris,
development, 513
host age, 513
parasitoid, 513
Pseudaletia unipuncta, 513
golf course,
Exomala orientalis, 14
temporal spatial distribution, 14
grape,
baculovirus, 868
enzyme-linked immunosorbent assay, 868
epizootiology, 868
microbial control, 868
vineyard, 868
grape phylloxera,
Daktolosphaira vitifoliae, 1004
distribution, 1004
extinction-colonization, 1004
population size, 1004
Vitis, 1004
grasshoppers,
Ageneotettix deorum, 347
hatch prediction, 347
models, 347
postdiapause development, 347
rangeland, 347
ground beetles,
biological control, 1145
conservation, 1145
habitat management, 1145
pitfall traps, 1145
refuge habitats, 1145

H

habitat management,
biological control, 1145
conservation, 1145
ground beetles, 1145
pitfall traps, 1145
refuge habitats, 1145
habitat structure,
Hogna helluo, 1052
Pardosa milvina, 1052
phenology, 1052
prey availability, 1052
tillage, 1052
habitats,
aphid parasitoid, 61
Diaeretiella rapae, 61
host range, 61
host switching, 61
Haematobia irritans,
diapause, 387
simulation model, 387
spring emergence, 387
temperature, 387
hatch prediction,
Ageneotettix deorum, 347
grasshoppers, 347
models, 347
postdiapause development, 347
rangeland, 347
heat shock,
acclimatization, 116
heat shock protein persistence, 116
thermotolerance, 116
Trichogramma brassicae, 116
heat shock protein persistence,
acclimatization, 116
heat shock, 116
thermotolerance, 116
Trichogramma brassicae, 116
Helianthus annuus,
Apis mellifera, 1172
Bacillus thuringiensis, 1172
Cochylis hospes, 1172
herbicide resistant germplasm,
delayed flood, 1130
Lissorhoptrus oryzophilus, 1130
rice plant, 1130
rice water weevil, 1130
herbivory, 30, 1044, 1136
Asclepias, 1044
defoliation, 1136
economic injury level, 1136
florivory, 1044
flowers, 1044
Glycine max, 1136

induced resistance, 30
insect injury, 1136
jasmonic acid, 30
leaf area index, 1136
plant defense, 30
plant–insect interactions, 30
Tetraopes, 1044
tomato, 30
Heterorhabditis,
entomopathogenic nematodes, 876
infectivity, 876
lepidopteran cotton pests, 876
Steinernema, 876
temperature, 876
Heterorhabditis bacteriophora, 128, 907, 1021
alfalfa snout beetle, 128
cadaver, 907
entomopathogenic nematode, 128
infectivity, 907
movement, 1021
porosity, 1021
soil compaction, 1021
Steinernema carpocapsae, 907, 1021
Steinernema glaseri, 1021
Hieracium pilosella,
Aulacidea pilosellae, 1
biological control, 1
phytophagous insect fauna, 1
Tephritis ruralis, 1
Hippodamia convergens, 518, 812
degree-days, 518
developmental threshold, 518
mortality, 518
Myzus persicae, 812
olfaction, 812
olfactometer, 812
scanning electron microscopy, 812
sensilla, 812
weights, 518
Hogna helluo,
habitat structure, 1052
Pardosa milvina, 1052
phenology, 1052
prey availability, 1052
tillage, 1052
horizontal toxicant transfer,
bait, 431
Blattella germanica, 431
coprophagy, 431
host acceptance, 137, 858
Anthonomus grandis, 137
Aphelinus asychis, 858
Aphidius matricariae, 858
biological control, 858
Callosobruchus maculatus, 137
Catolaccus grandis, 137
Diuraphis noxia, 858
factitious host, 137
host defense, 858
host age,
development, 513
Glyptapanteles militaris, 513
parasitoid, 513
Pseudaletia unipuncta, 513
host colonization,
host phenology, 461
local adaptation, 461
Operophtera brumata, 461
polyphagy, 461
host defense,
Aphelinus asychis, 858
Aphidius matricariae, 858
biological control, 858
Diuraphis noxia, 858
host acceptance, 858
host phenology,
host colonization, 461
local adaptation, 461
Operophtera brumata, 461
polyphagy, 461
host range,
aphid parasitoid, 61
Diaeretiella rapae, 61
habitats, 61
host switching, 61
host switching,
aphid parasitoid, 61
Diaeretiella rapae, 61
habitats, 61
host range, 61
host-finding,
attraction, 973
Colorado potato beetle, 973
Leptinotarsa decemlineata, 973
volicitin, 973

I

induced resistance,
herbivory, 30
jasmonic acid, 30
plant defense, 30
plant–insect interactions, 30

tomato, 30
induced response,
Bemisia argentifolii, 353
interspecific competition, 353
omnivory, 353
Trichoplusia ni, 353
inert dust,
biological control, 529
diatomaceous earth, 529
parasitoid, 529
stored products, 529
infection model,
Entomophaga maimaiga, 1162
infection reservoir, 1162
Lymantria dispar, 1162
rainfall, 1162
resting spores, 1162
infection reservoir,
Entomophaga maimaiga, 1162
infection model, 1162
Lymantria dispar, 1162
rainfall, 1162
resting spores, 1162
infectivity, 876, 907
cadaver, 907
entomopathogenic nematodes, 876
Heterorhabditis, 876
Heterorhabditis bacteriophora, 907
lepidopteran cotton pests, 876
Steinernema, 876
Steinernema carpocapsae, 907
temperature, 876
insect,
arthropod, 22
developmental rate, 22
model, 22
temperature, 22
insect coldhardiness,
Adelges tsugae, 1085
Tsuga canadensis, 1085
Tsuga caroliniana, 1085
insect injury,
defoliation, 1136
economic injury level, 1136
Glycine max, 1136
herbivory, 1136
leaf area index, 1136
insect pest management,
behavior, 1124
Coenosia tigrina, 1124
Delia radicum, 1124
exclusion fences, 1124
physical control, 1124
insect resistance,
Bacillus thuringiensis, 108
Chrysomela scripta, 108
cottonwood, 108
genetic engineering, 108
poplars, 108
insecticidal dusts,
body water content, 979
Callosobruchus maculatus, 979
cuticular permeability, 979
desiccants, 979
water loss, 979
insect–plant interactions,
conservation biological control, 1036
facultative phytophagy, 1036
flower bug, 1036
Orius insidiosus, 1036
integrated control, 836, 1060
Ephestia kuehniella, 1060
flour mills, 1060
population dynamics, 836, 1060
Prostephanus truncatus, 836
simulation model, 836, 1060
Sitophilus zeamais, 836
stored maize, 836
temperature, 1060
integrated pest management,
computer simulation, 505
population models, 505
interspecific competition,
Bemisia argentifolii, 353
induced response, 353
omnivory, 353
Trichoplusia ni, 353
interspecific intraguild predation,
cannibalism, 1092
coccinellids, 1092
dispersal, 1092
prey scarcity, 1092
interspecific predation,
cannibalism, 523
predaceous mites, 523
predation preference, 523
intrinsic rate of increase,
Bursaphelenchus xylophilus, 407
fertility table, 407
Monochamus carolinensis, 407

J

jasmonic acid,
herbivory, 30
induced resistance, 30
plant defense, 30
plant–insect interactions, 30
tomato, 30

L

lady beetle,
biological control, 901
Coccinellid, 901
feeding behavior, 901
predators, 901
whitefly, 901
large herbivorous dung,
dung beetle communities, 420
Mediterranean ecosystem, 420
rabbit dung, 420
Scarabaeoidea, 420
trophic resources, 420
Larra,
Apis, 367
Bombus, 367
Cerceris, 367
nontarget insects, 367
leaf area index,
defoliation, 1136
economic injury level, 1136
Glycine max, 1136
herbivory, 1136
insect injury, 1136
leafrollers,
apple, 81
mating disruption, 81
Pandemis, 81
sex pheromones, 81
Lepidoptera,
(E, E)-8,10-dodecadienol, 775
Cydia pomonella, 775
sex pheromone, 775
Tortricidae, 775
wind tunnel, 775
lepidopteran cotton pests,
entomopathogenic nematodes, 876
Heterorhabditis, 876
infectivity, 876
Steinernema, 876
temperature, 876
Leptinotarsa decemlineata,
attraction, 973
Colorado potato beetle, 973
host-finding, 973
volicitin, 973
lettuce,
aster yellows phytoplasma, 1101
gender, 1101
Macrosteles quadrilineatus, 1101
transmission, 1101
vector, 1101
life history parameters,
Bemisia tabaci, 831
melon, 831
resistance, 831
whitefly, 831
life table, 412, 895
brown citrus aphid, 895
citrus tristeza, 895
developmental rate, 412
population growth, 895
Toxoptera citricida, 412
vector, 895
virus, 895
Lissorhoptrus oryzophilus,
delayed flood, 1130
herbicide resistant germplasm, 1130
rice plant, 1130
rice water weevil, 1130
local adaptation,
host colonization, 461
host phenology, 461
Operophtera brumata, 461
polyphagy, 461
Lolium perenne,
Ataenius spretulus, 358
Bacillus popilliae, 358
mowing height, 358
predators, 358
turfgrass, 358
Lymantria dispar, 485, 1162
100-tree plots, 485
Entomophaga maimaiga, 485, 1162
fixed-radius plots, 485
infection model, 1162
infection reservoir, 1162
nuclear polyhedrosis virus, 485
rainfall, 1162
resting spores, 1162
timed walks, 485

M

Macrolepidoptera,
abundance, 795
sampling, 795

species richness, 795
traps, 795
weather, 795
Macrosteles quadrilineatus, 851, 1101
aster yellows phytoplasma, 1101
binomial sampling, 851
bootstrap simulation, 851
carrot, 851
gender, 1101
lettuce, 1101
resampling software, 851
transmission, 1101
vector, 1101
maize silk volatiles,
attractants, 961
Diabrotica barberi, 961
Diabrotica virgifera virgifera, 961
repellents, 961
semiochemicals, 961
Malacosoma californicum pluviale, 44, 1107
caterpillars, 1107
experimental epizootiology, 1107
forest Lepidoptera, 44
nuclear polyhedrosis virus, 44
nuclearpolyhedrovirus, 1107
population cycles, 44, 1107
population dynamics, 44
tent caterpillars, 44
virus transmission, 1107
management,
adult abundance, 398
Anthonomus signatus, 398
predictive model, 398
mark–recapture,
cannibalism, 823
Coptotermes acinaciformis, 823
mound-building, 823
Nile Blue A, 823
mark-recapture experiment,
bark beetles, 496
Dendroctonus frontalis, 496
dispersal, 496
pest management strategies, 496
spatial dynamics, 496
mass rearing,
biological control, 1183
Ceraeochrysa spp., 1183
development, 1183
reproduction, 1183
mating,
Colorado potato beetle, 425
flight, 425
resistance management, 425
transgenic plants, 425
mating behavior,
Ceratitis capitata, 780
trimedlure, 780
mating disruption,
apple, 81
leafrollers, 81
Pandemis, 81
sex pheromones, 81
mealybug,
buffalograss, 123
parasitoid, 123
Rhopus nigroclavatus, 123
Tridiscus sporoboli, 123
Medicago sativa,
alfalfa, 1067
foraging behavior, 1067
Megachile rotundata, 1067
pollination ecology, 1067
seed production, 1067
Mediterranean ecosystem,
dung beetle communities, 420
large herbivorous dung, 420
rabbit dung, 420
Scarabaeoidea, 420
trophic resources, 420
Megachile rotundata,
alfalfa, 1067
foraging behavior, 1067
Medicago sativa, 1067
pollination ecology, 1067
seed production, 1067
Megalurothrips sjostedti,
biological control, 1154
cowpea, 1154
Metarhizium anisopliae, 1154
synthetic insecticide nontarget organisms, 1154
thrips, 1154
melon,
Bemisia tabaci, 831
life history parameters, 831
resistance, 831
whitefly, 831
mesophyll-feeder,
azalea, 998
azalea lace bug, 998
development time, 998

nitrogen fertilizer, 998
survivorship, 998
Metarhizium anisopliae,
biological control, 1154
cowpea, 1154
Megalurothrips sjostedti, 1154
synthetic insecticide nontarget organisms, 1154
thrips, 1154
microbial control,
baculovirus, 868
enzyme-linked immunosorbent assay, 868
epizootiology, 868
grape, 868
vineyard, 868
model,
arthropod, 22
developmental rate, 22
insect, 22
temperature, 22
models,
Ageneotettix deorum, 347
grasshoppers, 347
hatch prediction, 347
postdiapause development, 347
rangeland, 347
Monochamus carolinensis,
Bursaphelenchus xylophilus, 407
fertility table, 407
intrinsic rate of increase, 407
mortality, 9, 439, 518, 935
degree-days, 518
dehydration, 935
developmental threshold, 518
emergence patterns, 9
Hippodamia convergens, 518
parasitoid, 9
Periplaneta americana, 935
Periplaneta fuliginosa, 935
Pieris rapae, 439
predation, 439
saturation deficit, 935
Shecophaga, 9
survival, 439
Vespula, 9
water relations, 935
weights, 518
yellowjackets, 9
mortality factors,
argan fruit, 53
Ceratitis capitata, 53
number of generations, 53
oviposition, 53
population dynamics, 53
mound-building,
cannibalism, 823
Coptotermes acinaciformis, 823
mark–recapture, 823
Nile Blue A, 823
mountain pine beetle,
climate change, 924
direct temperature control, 924
phenology, 924
seasonality, 924
temperature-dependent development, 924
movement,
Heterorhabditis bacteriophora, 1021
porosity, 1021
soil compaction, 1021
Steinernema carpocapsae, 1021
Steinernema glaseri, 1021
mowing height,
Ataenius spretulus, 358
Bacillus popilliae, 358
Lolium perenne, 358
predators, 358
turfgrass, 358
Myzus persicae, 812, 983
alarm pheromone, 983
aphid, 983
Diuraphis noxia, 983
Hippodamia convergens, 812
olfaction, 812
olfactometer, 812
Paecilomyces fumosoroseus, 983
scanning electron microscopy, 812
semiochemicals, 983
sensilla, 812

N

natural enemies,
Beauveria bassiana, 863
rangeland grasshoppers, 863
Neodiprion gillettei,
above-below-ground growth, 38
compensatory growth, 38
Pinus ponderosa, 38
sawfly defoliation, 38
Nilaparvata lugens,
biological control, 1177

enzyme-linked immunosorbent assay, 1177
predation, 1177
spider, 1177
Nile Blue A,
cannibalism, 823
Coptotermes acinaciformis, 823
mark–recapture, 823
mound-building, 823
nitrogen fertilizer,
azalea, 998
azalea lace bug, 998
development time, 998
mesophyll-feeder, 998
survivorship, 998
nontarget insects,
Apis, 367
Bombus, 367
Cerceris, 367
Larra, 367
nuclear polyhedrosis virus, 44, 485
100-tree plots, 485
Entomophaga maimaiga, 485
fixed-radius plots, 485
forest Lepidoptera, 44
Lymantria dispar, 485
Malacosoma californicum pluviale, 44
population cycles, 44
population dynamics, 44
tent caterpillars, 44
timed walks, 485
nuclearpolyhedrovirus,
caterpillars, 1107
experimental epizootiology, 1107
Malacosoma californicum pluviale, 1107
population cycles, 1107
virus transmission, 1107
number of generations,
argan fruit, 53
Ceratitis capitata, 53
mortality factors, 53
oviposition, 53
population dynamics, 53
nymphal survivorship,
feeding injury, 456
Freeman maple, 456
oviposition, 456
potato leafhopper, 456
red maple, 456

O

odor quantification,
olfactory stimulation, 947
pheromones, 947
plant volatiles, 947
release rate, 947
ultrasound, 947
olfaction,
Hippodamia convergens, 812
Myzus persicae, 812
olfactometer, 812
scanning electron microscopy, 812
sensilla, 812
olfactometer,
Hippodamia convergens, 812
Myzus persicae, 812
olfaction, 812
scanning electron microscopy, 812
sensilla, 812
olfactory stimulation,
odor quantification, 947
pheromones, 947
plant volatiles, 947
release rate, 947
ultrasound, 947
olfactory stimuli,
Bactrocera dorsalis, 1140
pest sampling, 1140
protein, 1140
visual stimuli, 1140
omnivory,
Bemisia argentifolii, 353
induced response, 353
interspecific competition, 353
Trichoplusia ni, 353
100-tree plots,
Entomophaga maimaiga, 485
fixed-radius plots, 485
Lymantria dispar, 485
nuclear polyhedrosis virus, 485
timed walks, 485
Operophtera brumata, 96, 461
biological control, 96
Cyzenis albicans, 96
host colonization, 461
host phenology, 461
local adaptation, 461
polyphagy, 461
simultaneous introduction, 96
winter moth, 96
Orius insidiosus,

conservation biological control, 1036
facultative phytophagy, 1036
flower bug, 1036
insect–plant interactions, 1036
oviposition, 53, 456
argan fruit, 53
Ceratitis capitata, 53
feeding injury, 456
Freeman maple, 456
mortality factors, 53
number of generations, 53
nymphal survivorship, 456
population dynamics, 53
potato leafhopper, 456
red maple, 456

P

Paecilomyces fumosoroseus,
alarm pheromone, 983
aphid, 983
Diuraphis noxia, 983
Myzus persicae, 983
semiochemicals, 983
Pandemis,
apple, 81
leafrollers, 81
mating disruption, 81
sex pheromones, 81
Panonychus ulmi,
Amblyseius fallacis, 72
biological control, 72
encapsulated cypermethrin, 72
parasitic ants,
biological control, 88
colony composition, 88
fire ants, 88
Solenopsis daguerrei, 88
Solenopsis richteri, 88
parasitoid, 9, 123, 513, 529
biological control, 529
buffalograss, 123
development, 513
diatomaceous earth, 529
emergence patterns, 9
Glyptapanteles militaris, 513
host age, 513
inert dust, 529
mealybug, 123
mortality, 9
Pseudaletia unipuncta, 513
Rhopus nigroclavatus, 123
Shecophaga, 9
stored products, 529
Tridiscus sporoboli, 123
Vespula, 9
yellowjackets, 9
parasitoids,
community structure, 1114
density dependence, 1114
Diadegma psilocorsis, 1114
Psilocorsis quercicella, 1114
Quercus alba, 1114
Pardosa milvina,
habitat structure, 1052
Hogna helluo, 1052
phenology, 1052
prey availability, 1052
tillage, 1052
peas,
aggregation, 477
Sitona lineatus, 477
spatiotemporal analysis, 477
tillage, 477
Pectinophora gossypiella,
biological invasions, 445
climate matching software, 445
cotton, 445
establishment, 445
risk assessment, 445
Periplaneta americana,
dehydration, 935
mortality, 935
Periplaneta fuliginosa, 935
saturation deficit, 935
water relations, 935
Periplaneta fuliginosa,
dehydration, 935
mortality, 935
Periplaneta americana, 935
saturation deficit, 935
water relations, 935
pest management strategies,
bark beetles, 496
Dendroctonus frontalis, 496
dispersal, 496
mark-recapture experiment, 496
spatial dynamics, 496
pest sampling,
Bactrocera dorsalis, 1140
olfactory stimuli, 1140
protein, 1140
visual stimuli, 1140

phenology, 924, 1052
climate change, 924
direct temperature control, 924
habitat structure, 1052
Hogna helluo, 1052
mountain pine beetle, 924
Pardosa milvina, 1052
prey availability, 1052
seasonality, 924
temperature-dependent development, 924
tillage, 1052
pheromone,
female behavior, 943
pheromone autodetection, 943
Vitacea polistiformis, 943
pheromone autodetection,
female behavior, 943
pheromone, 943
Vitacea polistiformis, 943
pheromones,
odor quantification, 947
olfactory stimulation, 947
plant volatiles, 947
release rate, 947
ultrasound, 947
phospholipids,
cold heat tolerance, 968
Drosophila, 968
energy storage, 968
triacylglycerols, 968
photosynthesis,
antibiosis, 787
chlorophyll fluorescence, 787
Russian wheat aphid, 787
tolerance, 787
wheat, 787
phylloxera,
biological control, 890
Daktulosphaira vitifolia, 890
entomopathogenic nematodes, 890
soil moisture, 890
Vitis, 890
physical control,
behavior, 1124
Coenosia tigrina, 1124
Delia radicum, 1124
exclusion fences, 1124
insect pest management, 1124
phytophagous insect fauna,
Aulacidea pilosellae, 1
biological control, 1
Hieracium pilosella, 1
Tephritis ruralis, 1
Pieris rapae,
mortality, 439
predation, 439
survival, 439
pine wilt disease,
Bursaphelenchus xylophilus, 1014
flight, 1014
flight mill, 1014
pinewood nematode, 1014
pinewood nematode,
Bursaphelenchus xylophilus, 1014
flight, 1014
flight mill, 1014
pine wilt disease, 1014
Pinus ponderosa,
above-below-ground growth, 38
compensatory growth, 38
Neodiprion gillettei, 38
sawfly defoliation, 38
pitfall traps,
biological control, 1145
conservation, 1145
ground beetles, 1145
habitat management, 1145
refuge habitats, 1145
plant defense,
herbivory, 30
induced resistance, 30
jasmonic acid, 30
plant–insect interactions, 30
tomato, 30
plant volatiles,
odor quantification, 947
olfactory stimulation, 947
pheromones, 947
release rate, 947
ultrasound, 947
plant–insect interactions,
herbivory, 30
induced resistance, 30
jasmonic acid, 30
plant defense, 30
tomato, 30
pollination ecology,
alfalfa, 1067
foraging behavior, 1067
Medicago sativa, 1067
Megachile rotundata, 1067

seed production, 1067
polyphagy,
host colonization, 461
host phenology, 461
local adaptation, 461
Operophtera brumata, 461
poplars,
Bacillus thuringiensis, 108
Chrysomela scripta, 108
cottonwood, 108
genetic engineering, 108
insect resistance, 108
population cycles, 44, 1107
caterpillars, 1107
experimental epizootiology, 1107
forest Lepidoptera, 44
Malacosoma californicum pluviale, 44, 1107
nuclear polyhedrosis virus, 44
nuclearpolyhedrovirus, 1107
population dynamics, 44
tent caterpillars, 44
virus transmission, 1107
population dynamics, 44, 53, 372, 836, 1060
argan fruit, 53
Ceratitis capitata, 53
dispersal, 372
Ephestia kuehniella, 1060
flour mills, 1060
forest Lepidoptera, 44
integrated control, 836, 1060
Malacosoma californicum pluviale, 44
mortality factors, 53
nuclear polyhedrosis virus, 44
number of generations, 53
oviposition, 53
population cycles, 44
Prosapia nr *bicincta*, 372
Prosapia simulans, 372
Prostephanus truncatus, 836
simulation model, 836, 1060
Sitophilus zeamais, 836
stored maize, 836
temperature, 1060
tent caterpillars, 44
population growth,
brown citrus aphid, 895
citrus tristeza, 895
life table, 895
vector, 895
virus, 895
population models,
computer simulation, 505
integrated pest management, 505
population size,
Daktolosphaira vitifoliae, 1004
distribution, 1004
extinction-colonization, 1004
grape phylloxera, 1004
Vitis, 1004
porosity,
Heterorhabditis bacteriophora, 1021
movement, 1021
soil compaction, 1021
Steinernema carpocapsae, 1021
Steinernema glaseri, 1021
postdiapause development,
Ageneotettix deorum, 347
grasshoppers, 347
hatch prediction, 347
models, 347
rangeland, 347
potato leafhopper,
feeding injury, 456
Freeman maple, 456
nymphal survivorship, 456
oviposition, 456
red maple, 456
predaceous mites,
cannibalism, 523
interspecific predation, 523
predation preference, 523
predation, 439, 1177
biological control, 1177
enzyme-linked immunosorbent assay, 1177
mortality, 439
Nilaparvata lugens, 1177
Pieris rapae, 439
spider, 1177
survival, 439
predation preference,
cannibalism, 523
interspecific predation, 523
predaceous mites, 523
predators, 358, 901
Ataenius spretulus, 358
Bacillus popilliae, 358
biological control, 901
Coccinellid, 901

feeding behavior, 901
lady beetle, 901
Lolium perenne, 358
mowing height, 358
turfgrass, 358
whitefly, 901
predictive model,
adult abundance, 398
Anthonomus signatus, 398
management, 398
prey availability,
habitat structure, 1052
Hogna helluo, 1052
Pardosa milvina, 1052
phenology, 1052
tillage, 1052
prey scarcity,
cannibalism, 1092
coccinellids, 1092
dispersal, 1092
interspecific intraguild predation, 1092
Prosapia nr *bicincta*,
dispersal, 372
population dynamics, 372
Prosapia simulans, 372
Prosapia simulans,
dispersal, 372
population dynamics, 372
Prosapia nr *bicincta*, 372
Prostephanus truncatus,
integrated control, 836
population dynamics, 836
simulation model, 836
Sitophilus zeamais, 836
stored maize, 836
protein,
Bactrocera dorsalis, 1140
olfactory stimuli, 1140
pest sampling, 1140
visual stimuli, 1140
Pseudaletia unipuncta,
development, 513
Glyptapanteles militaris, 513
host age, 513
parasitoid, 513
Psilocorsis quercicella,
community structure, 1114
density dependence, 1114
Diadegma psilocorsis, 1114
parasitoids, 1114
Quercus alba, 1114

Q

Quercus alba,
community structure, 1114
density dependence, 1114
Diadegma psilocorsis, 1114
parasitoids, 1114
Psilocorsis quercicella, 1114

R

rabbit dung,
dung beetle communities, 420
large herbivorous dung, 420
Mediterranean ecosystem, 420
Scarabaeoidea, 420
trophic resources, 420
rainfall,
Entomophaga maimaiga, 1162
infection model, 1162
infection reservoir, 1162
Lymantria dispar, 1162
resting spores, 1162
rangeland,
Ageneotettix deorum, 347
grasshoppers, 347
hatch prediction, 347
models, 347
postdiapause development, 347
rangeland grasshoppers,
Beauveria bassiana, 863
natural enemies, 863
red maple,
feeding injury, 456
Freeman maple, 456
nymphal survivorship, 456
oviposition, 456
potato leafhopper, 456
refuge habitats,
biological control, 1145
conservation, 1145
ground beetles, 1145
habitat management, 1145
pitfall traps, 1145
release rate,
odor quantification, 947
olfactory stimulation, 947
pheromones, 947
plant volatiles, 947
ultrasound, 947
repellent,

arrestant, 954
attractant, 954
Cydia pomonella, 954
deterrent, 954
repellents,
attractants, 961
Diabrotica barberi, 961
Diabrotica virgifera virgifera, 961
maize silk volatiles, 961
semiochemicals, 961
reproduction,
biological control, 1183
Ceraeochrysa spp., 1183
development, 1183
mass rearing, 1183
resampling software,
binomial sampling, 851
bootstrap simulation, 851
carrot, 851
Macrosteles quadrilineatus, 851
resistance,
Bemisia tabaci, 831
life history parameters, 831
melon, 831
whitefly, 831
resistance management,
Colorado potato beetle, 425
flight, 425
mating, 425
transgenic plants, 425
resting spores,
Entomophaga maimaiga, 1162
infection model, 1162
infection reservoir, 1162
Lymantria dispar, 1162
rainfall, 1162
Rhopus nigroclavatus,
buffalograss, 123
mealybug, 123
parasitoid, 123
Tridiscus sporoboli, 123
rice plant,
delayed flood, 1130
herbicide resistant germplasm, 1130
Lissorhoptrus oryzophilus, 1130
rice water weevil, 1130
rice water weevil,
delayed flood, 1130
herbicide resistant germplasm, 1130
Lissorhoptrus oryzophilus, 1130
rice plant, 1130
risk assessment,
biological invasions, 445
climate matching software, 445
cotton, 445
establishment, 445
Pectinophora gossypiella, 445
Russian wheat aphid,
antibiosis, 787
chlorophyll fluorescence, 787
photosynthesis, 787
tolerance, 787
wheat, 787

S

sampling,
abundance, 795
Macrolepidoptera, 795
species richness, 795
traps, 795
weather, 795
saturation deficit,
dehydration, 935
mortality, 935
Periplaneta americana, 935
Periplaneta fuliginosa, 935
water relations, 935
sawfly defoliation,
above-below-ground growth, 38
compensatory growth, 38
Neodiprion gillettei, 38
Pinus ponderosa, 38
scanning electron microscopy,
Hippodamia convergens, 812
Myzus persicae, 812
olfaction, 812
olfactometer, 812
sensilla, 812
Scarabaeoidea,
dung beetle communities, 420
large herbivorous dung, 420
Mediterranean ecosystem, 420
rabbit dung, 420
trophic resources, 420
seasonality,
climate change, 924
direct temperature control, 924
mountain pine beetle, 924
phenology, 924
temperature-dependent development, 924
seed production,
alfalfa, 1067

foraging behavior, 1067
Medicago sativa, 1067
Megachile rotundata, 1067
pollination ecology, 1067
semiochemicals, 961, 983
alarm pheromone, 983
aphid, 983
attractants, 961
Diabrotica barberi, 961
Diabrotica virgifera virgifera, 961
Diuraphis noxia, 983
maize silk volatiles, 961
Myzus persicae, 983
Paecilomyces fumosoroseus, 983
repellents, 961
sensilla,
Hippodamia convergens, 812
Myzus persicae, 812
olfaction, 812
olfactometer, 812
scanning electron microscopy, 812
sex pheromone,
(E, E)-8,10-dodecadienol, 775
Cydia pomonella, 775
Lepidoptera, 775
Tortricidae, 775
wind tunnel, 775
sex pheromones,
apple, 81
leafrollers, 81
mating disruption, 81
Pandemis, 81
Shecophaga,
emergence patterns, 9
mortality, 9
parasitoid, 9
Vespula, 9
yellowjackets, 9
simulation model, 387, 836, 1060
diapause, 387
Ephestia kuehniella, 1060
flour mills, 1060
Haematobia irritans, 387
integrated control, 836, 1060
population dynamics, 836, 1060
Prostephanus truncatus, 836
Sitophilus zeamais, 836
spring emergence, 387
stored maize, 836
temperature, 387, 1060
simultaneous introduction,
biological control, 96
Cyzenis albicans, 96
Operophtera brumata, 96
winter moth, 96
Sitona lineatus,
aggregation, 477
peas, 477
spatiotemporal analysis, 477
tillage, 477
Sitophilus zeamais,
integrated control, 836
population dynamics, 836
Prostephanus truncatus, 836
simulation model, 836
stored maize, 836
soil compaction,
Heterorhabditis bacteriophora, 1021
movement, 1021
porosity, 1021
Steinernema carpocapsae, 1021
Steinernema glaseri, 1021
soil moisture,
biological control, 890
Daktulosphaira vitifolia, 890
entomopathogenic nematodes, 890
phylloxera, 890
Vitis, 890
Solenopsis daguerrei,
biological control, 88
colony composition, 88
fire ants, 88
parasitic ants, 88
Solenopsis richteri, 88
Solenopsis richteri,
biological control, 88
colony composition, 88
fire ants, 88
parasitic ants, 88
Solenopsis daguerrei, 88
spatial dynamics,
bark beetles, 496
Dendroctonus frontalis, 496
dispersal, 496
mark-recapture experiment, 496
pest management strategies, 496
spatiotemporal analysis,
aggregation, 477
peas, 477
Sitona lineatus, 477
tillage, 477
species richness, 795, 986

abundance, 795
biodiversity, 986
Braconidae, 986
complementarity, 986
forest management, 986
Macrolepidoptera, 795
sampling, 795
traps, 795
weather, 795
spider,
biological control, 1177
enzyme-linked immunosorbent assay, 1177
Nilaparvata lugens, 1177
predation, 1177
spider mites,
biological control, 915
discriminant analysis, 915
generalist-specialist predators, 915
genetic algorithm, 915
Tetranychidae, 915
spring emergence,
diapause, 387
Haematobia irritans, 387
simulation model, 387
temperature, 387
Steinernema,
entomopathogenic nematodes, 876
Heterorhabditis, 876
infectivity, 876
lepidopteran cotton pests, 876
temperature, 876
Steinernema carpocapsae, 907, 1021
cadaver, 907
Heterorhabditis bacteriophora, 907, 1021
infectivity, 907
movement, 1021
porosity, 1021
soil compaction, 1021
Steinernema glaseri, 1021
Steinernema glaseri,
Heterorhabditis bacteriophora, 1021
movement, 1021
porosity, 1021
soil compaction, 1021
Steinernema carpocapsae, 1021
stored maize,
integrated control, 836
population dynamics, 836
Prostephanus truncatus, 836
simulation model, 836
Sitophilus zeamais, 836
stored products,
biological control, 529
diatomaceous earth, 529
inert dust, 529
parasitoid, 529
subterranean termites,
feeding behavior, 1077
foraging behavior, 1077
wood consumption, 1077
survival,
mortality, 439
Pieris rapae, 439
predation, 439
survivorship,
azalea, 998
azalea lace bug, 998
development time, 998
mesophyll-feeder, 998
nitrogen fertilizer, 998
synthetic insecticide nontarget organisms,
biological control, 1154
cowpea, 1154
Megalurothrips sjostedti, 1154
Metarhizium anisopliae, 1154
thrips, 1154

T

temperature, 22, 387, 876, 1060
arthropod, 22
developmental rate, 22
diapause, 387
entomopathogenic nematodes, 876
Ephestia kuehniella, 1060
flour mills, 1060
Haematobia irritans, 387
Heterorhabditis, 876
infectivity, 876
insect, 22
integrated control, 1060
lepidopteran cotton pests, 876
model, 22
population dynamics, 1060
simulation model, 387, 1060
spring emergence, 387
Steinernema, 876
temperature-dependent development,
climate change, 924
direct temperature control, 924
mountain pine beetle, 924

phenology, 924
seasonality, 924
temporal spatial distribution,
Exomala orientalis, 14
golf course, 14
tent caterpillars,
forest Lepidoptera, 44
Malacosoma californicum pluviale, 44
nuclear polyhedrosis virus, 44
population cycles, 44
population dynamics, 44
Tephritis ruralis,
Aulacidea pilosellae, 1
biological control, 1
Hieracium pilosella, 1
phytophagous insect fauna, 1
Tetranychidae,
biological control, 915
discriminant analysis, 915
generalist-specialist predators, 915
genetic algorithm, 915
spider mites, 915
Tetraopes,
Asclepias, 1044
florivory, 1044
flowers, 1044
herbivory, 1044
thermotolerance,
acclimatization, 116
heat shock, 116
heat shock protein persistence, 116
Trichogramma brassicae, 116
thrips,
biological control, 1154
cowpea, 1154
Megalurothrips sjostedti, 1154
Metarhizium anisopliae, 1154
synthetic insecticide nontarget organisms, 1154
tillage, 477, 1052
aggregation, 477
habitat structure, 1052
Hogna helluo, 1052
Pardosa milvina, 1052
peas, 477
phenology, 1052
prey availability, 1052
Sitona lineatus, 477
spatiotemporal analysis, 477
timed walks,
100-tree plots, 485
Entomophaga maimaiga, 485
fixed-radius plots, 485
Lymantria dispar, 485
nuclear polyhedrosis virus, 485
tolerance,
antibiosis, 787
chlorophyll fluorescence, 787
photosynthesis, 787
Russian wheat aphid, 787
wheat, 787
tomato,
herbivory, 30
induced resistance, 30
jasmonic acid, 30
plant defense, 30
plant–insect interactions, 30
Tortricidae,
(E, E)-8,10-dodecadienol, 775
Cydia pomonella, 775
Lepidoptera, 775
sex pheromone, 775
wind tunnel, 775
Toxoptera citricida,
developmental rate, 412
life table, 412
transgenic plants,
Colorado potato beetle, 425
flight, 425
mating, 425
resistance management, 425
transmission,
aster yellows phytoplasma, 1101
gender, 1101
lettuce, 1101
Macrosteles quadrilineatus, 1101
vector, 1101
traps,
abundance, 795
Macrolepidoptera, 795
sampling, 795
species richness, 795
weather, 795
triacylglycerols,
cold heat tolerance, 968
Drosophila, 968
energy storage, 968
phospholipids, 968
Trichogramma brassicae,
acclimatization, 116
heat shock, 116

heat shock protein persistence, 116
thermotolerance, 116
Trichoplusia ni,
Bemisia argentifolii, 353
induced response, 353
interspecific competition, 353
omnivory, 353
Tridiscus sporoboli,
buffalograss, 123
mealybug, 123
parasitoid, 123
Rhopus nigroclavatus, 123
trimedlure,
Ceratitis capitata, 780
mating behavior, 780
Trioxys curvicaudus,
canopy distribution, 845
Eucallipterus tiliae, 845
trophic resources,
dung beetle communities, 420
large herbivorous dung, 420
Mediterranean ecosystem, 420
rabbit dung, 420
Scarabaeoidea, 420
Tsuga canadensis,
Adelges tsugae, 1085
insect coldhardiness, 1085
Tsuga caroliniana, 1085
Tsuga caroliniana,
Adelges tsugae, 1085
insect coldhardiness, 1085
Tsuga canadensis, 1085
turfgrass,
Ataenius spretulus, 358
Bacillus popilliae, 358
Lolium perenne, 358
mowing height, 358
predators, 358

U

ultrasound,
odor quantification, 947
olfactory stimulation, 947
pheromones, 947
plant volatiles, 947
release rate, 947

V

vector, 895, 1101
aster yellows phytoplasma, 1101
brown citrus aphid, 895
citrus tristeza, 895
gender, 1101
lettuce, 1101
life table, 895
Macrosteles quadrilineatus, 1101
population growth, 895
transmission, 1101
virus, 895
Vespula,
emergence patterns, 9
mortality, 9
parasitoid, 9
Shecophaga, 9
yellowjackets, 9
vineyard,
baculovirus, 868
enzyme-linked immunosorbent assay, 868
epizootiology, 868
grape, 868
microbial control, 868
virus,
brown citrus aphid, 895
citrus tristeza, 895
life table, 895
population growth, 895
vector, 895
virus transmission,
caterpillars, 1107
experimental epizootiology, 1107
Malacosoma californicum pluviale, 1107
nuclearpolyhedrovirus, 1107
population cycles, 1107
visual stimuli,
Bactrocera dorsalis, 1140
olfactory stimuli, 1140
pest sampling, 1140
protein, 1140
Vitacea polistiformis,
female behavior, 943
pheromone, 943
pheromone autodetection, 943
Vitis, 890, 1004
biological control, 890
Daktolosphaira vitifoliae, 1004
Daktulosphaira vitifolia, 890
distribution, 1004
entomopathogenic nematodes, 890
extinction-colonization, 1004
grape phylloxera, 1004

phylloxera, 890
population size, 1004
soil moisture, 890
volicitin,
attraction, 973
Colorado potato beetle, 973
host-finding, 973
Leptinotarsa decemlineata, 973

W

water loss,
body water content, 979
Callosobruchus maculatus, 979
cuticular permeability, 979
desiccants, 979
insecticidal dusts, 979
water relations,
dehydration, 935
mortality, 935
Periplaneta americana, 935
Periplaneta fuliginosa, 935
saturation deficit, 935
weather,
abundance, 795
Macrolepidoptera, 795
sampling, 795
species richness, 795
traps, 795
weights,
degree-days, 518
developmental threshold, 518
Hippodamia convergens, 518
mortality, 518
wheat,
antibiosis, 787
chlorophyll fluorescence, 787
photosynthesis, 787
Russian wheat aphid, 787
tolerance, 787
whitefly, 831, 901
Bemisia tabaci, 831
biological control, 901
Coccinellid, 901
feeding behavior, 901
lady beetle, 901
life history parameters, 831
melon, 831
predators, 901
resistance, 831
wind tunnel,
(E, E)-8,10-dodecadienol, 775
Cydia pomonella, 775
Lepidoptera, 775
sex pheromone, 775
Tortricidae, 775
winter moth,
biological control, 96
Cyzenis albicans, 96
Operophtera brumata, 96
simultaneous introduction, 96
wood consumption,
feeding behavior, 1077
foraging behavior, 1077
subterranean termites, 1077

Y

yellowjackets,
emergence patterns, 9
mortality, 9
parasitoid, 9
Shecophaga, 9
Vespula, 9